


Dihydrobotrydial

Farnesyl pyrophosphate was incorporated into dihydrobotrydial to the extent of 0.33% thereby establishing its sesquiterpenoid origin. By dissecting out possible isoprene units, several possible ways (fig.1 a - d) of forming the carbon skeleton can be envisaged.



The carbon skeleton of farnesyl pyrophosphate will be labelled from $[1-^{13}\text{C}]$ -acetate as shown in fig. 2(a) whilst $[1,2-^{13}\text{C}_2]$ -acetate will generate the coupling pattern as shown in fig. 2(b) provided it is fed at such a dilution that individual molecules of farnesyl pyrophosphate are not biosynthesized from more than one labelled acetate unit (vide infra). $[4,5-^{13}\text{C}_2]$ -Mevalonate generates the much simpler coupling pattern (fig. 2(c)).




figure 2

Both $[1,2-^{13}\text{C}_2]$ -acetate and $[4,5-^{13}\text{C}_2]$ -mevalonate were fed to Botrytis cinerea. Analysis of the ^{13}C n.m.r. spectra of the resultant dihydrobotrydial revealed coupling patterns which were consistent with either foldings (c) or (d). These foldings were distinguished by the use of $[1-^{13}\text{C}]$ -acetate. In the biosynthesis of polyketides, small (in intensity) couplings have sometimes been observed between acetate units. These arise by the multiple-labelling of the polyketide chain. Whilst studying the biosynthesis of the sesquiterpenoid trichothecin by Trichothecium roseum, we showed⁸ that feeding large quantities of acetate inhibited the formation of endogenous acetate by, for example, glycolysis. Hence if a pulse of $[1-^{13}\text{C}]$ -acetate is fed to the fungus, a situation arises in which, for a short time, the farnesyl pyrophosphate is made primarily from exogenous material and is multiply-labelled. If a bond is then created in a subsequent cyclization or rearrangement between two C-1 labelled centres, a new $^{13}\text{C}-^{13}\text{C}$ coupling is observed. Folding (c) will generate a coupling between C-8 and C-9 in dihydrobotrydial whilst folding (d) will generate couplings between C-6,C-7 and C-8. In practice the latter set of induced couplings were observed. This leads us to propose the following biosynthetic pathway (figure 3) which is reminiscent of that leading to the sesquiterpenoid hydrocarbon, caryophyllene.⁹




figure 3

The second phase of the problem involved the stereochemical fate of the mevalonoid hydrogen atoms. $[2\text{-}^3\text{H}_2, 2\text{-}^{14}\text{C}]$ -, $[2\text{-}(\text{R})\text{-}2\text{-}^3\text{H}, 2\text{-}^{14}\text{C}]$ -, $[4\text{-}(\text{R})\text{-}4\text{-}^3\text{H}, 2\text{-}^{14}\text{C}]$ -, $[5\text{-}^3\text{H}_2, 2\text{-}^{14}\text{C}]$ - and $[5\text{-}(\text{R})\text{-}5\text{-}^3\text{H}, 2\text{-}^{14}\text{C}]$ -Mevalonates were each fed to the fungus and the atom ratios obtained in the product are given in table 1

Table 1 Incorporation of Mevalonates into Dihydrobotrydial (1)

Mevalonate	$2\text{-}^3\text{H}, 2\text{-}^{14}\text{C}$	$4\text{-}(\text{R})\text{-}4\text{-}^3\text{H}, 2\text{-}^{14}\text{C}$	$5\text{-}^3\text{H}, 2\text{-}^{14}\text{C}$
$^3\text{H}:^{14}\text{C}$ atom ratio in (1)	4.9:3 (0.14%)		3.9:3 (0.52%)
$^3\text{H}:^{14}\text{C}$ atom ratio from stereospecific(R) MVA in (1)	2.86:3 (1.28%)	2.9:3 (0.2%)	3.3:3 (0.7%)
incorporation (%)			

All three $4(\text{R})\text{-}4\text{-}^3\text{H}$ mevalonoid hydrogen labels were incorporated. In dihydrobotrydial one of the centres (C-9) labelled by C-4 of mevalonate, is fully substituted and hence the corresponding hydrogen atom has been involved in a rearrangement. In the proposed biosynthetic scheme (fig.3), the rearrangement leads to a hydrogen atom appearing at C-2 thus generating the secondary methyl group. The rearrangement could take the form of two 1,2-shifts ($\text{H-9} \rightarrow \text{C-1}$; $\text{H-1} \rightarrow \text{C-2}$) or a direct 1,3-shift ($\text{H-9} \rightarrow \text{C-2}$) (see figure 4):





figure 4

These have been distinguished by both ^{13}C and ^2H n.m.r. methods. [$4\text{-}^2\text{H}_2, 4\text{-}^{13}\text{C}$] Mevalonic acid was prepared so that each carbon-13 label also bore a deuterium atom. The mevalonate was fed at such a dilution to *Botrytis cinerea* that the individual molecules of dihydrobotrydial contained only one labelled unit. In pathway (a) two deuterium atoms become detached from their carbon-13 partners whilst in pathway (b) only one atom becomes detached. This has consequences in both the carbon and deuterium n.m.r. spectra. In the proton noise decoupled ^{13}C n.m.r. spectrum, the $^1\text{H}:^{13}\text{C}$ couplings are removed and the relaxation mode of the carbon-13 is also modified. Both these effects lead to an enhancement of the signal. If deuterium with a spin of 1, is attached to the carbon atom, not only does the $^2\text{H}:^{13}\text{C}$ coupling remain in the proton noise decoupled spectrum but also the nuclear Overhauser enhancement is absent. Hence when deuterium is attached to a carbon-13 any biosynthetic enrichment shown by signal increase over the natural abundance spectrum, is quenched. Pathway (a) moves two deuterium atoms away from their carbon-13 partners and thus two out of the three enrichments will be revealed and only one will be quenched. On the other hand pathway (b) moves only one deuterium atom and hence only one enrichment will be revealed, that of the tertiary alcohol C-9. Since carbon-13 has a spin of $\frac{1}{2}$ whilst deuterium has a spin of 1, any $^{13}\text{C}-^2\text{H}$ heteronuclear coupling is more easily observed in the deuterium spectrum. Pathway (a) destroys two couplings and will afford two singlets and one doublet in the deuterium spectrum whilst pathway (b) gives one singlet and two doublets. Furthermore since the proton n.m.r. spectrum has been assigned, the deuterium chemical shifts indicates the sites of the labels. The carbon-13 n.m.r. results supported pathway (b) with only C-9 showing an enrichment. The deuterium n.m.r. spectrum of botryalolic acid ester (3), a relative of dihydrobotrydial and also produced by the fungus, was more clearly resolved. The $^2\text{H}-1$ signal and the $^2\text{H}-5$ signal appeared as doublets (J 19.5 and 21 Hz) whilst the $^2\text{H}-2$ signal appeared as a singlet each superimposed on the natural abundance spectrum. Hence there is a 1,3-hydride shift in dihydrobotrydial biosynthesis which is stereochemically much more acceptable producing an inversion of configuration at C-9, i.e. the hydroxyl group lies on the opposite face of the molecule to H-2 (see figure 5).



3

figure 5

The 2-(R)-2- ^3H - and 5-(R)-5- ^3H -mevalonate results provided some information on the formation of the hemi-acetal ring. Oxidation of the dihydrobotrydial to the corresponding lactone showed that the C-10 hydrogen atom was derived from the pro-5(R) position of mevalonic acid whilst degradation to the 10,15-dicarboxylic acid ester showed that C-15 bore only one 2- ^3H -mevalonid label and that this was a pro-2(R)-hydrogen atom. The distinction between the two enantiotopic hydrogen atoms at C-15 was made using deuterium labelling. The two hydrogen atoms may be distinguished in the ^1H n.m.r. spectrum by solvent and shift reagent studies. [$2\text{-}^2\text{H}_2$]-Acetate labels the C-2 position of MVA and hence C-15 in dihydrobotrydial. Examination of the deuterium n.m.r. spectrum of the labelled dihydrobotrydial derived from [$2\text{-}^2\text{H}_2$]-acetate showed that the mevalonid label was on the lower field (δ 4.20) of the two protons on the same face of the molecule as the two hydroxyl groups. The stereochemistry of labelling of farnesyl pyrophosphate at C-4 and C-5 from pro-2(R)- and pro-5(R) mevalonate is known. Consequently the formation of the hemi-acetal ring proceeds with overall retention of configuration at both centres (see figure 6). Although a number of mechanisms may be proposed to account for this, the dialdehyde, botrydial (4) was incorporated into dihydrobotrydial (1) to the extent of 32% whereas the reverse reaction proceeded in 1.09% yield. Thus the cleavage of the 10:15 bond in the formation of the hemi-acetal probably occurs through the dialdehyde which in turn could be formed via a trans-15 α ,10 β -glycol.




figure 6

The oxidation of a *trans*-glycol would lead to a dialdehyde. The subsequent reduction of the aldehyde at C-15 proceeds with the *re*-stereospecificity which is typical of a microbial dehydrogenase. A similar cleavage may occur in the formation of another sesquiterpenoid fungal metabolite, helminthosporal.

Alliacolide

The crystallographic data for the second fungal metabolite, alliacolide (2), did not permit the assignment of an absolute configuration. Degradation and circular dichroism measurements on the following compounds (see figure 7) have enabled us to define this.




figure 7

Examination of the culture broth of *Marasmius alliaceus* has led to the isolation of the following metabolites (see figure 8). Careful examination of the ^{13}C n.m.r. spectra of these and of a number of derivatives facilitated the assignment of the carbon-13 resonances whilst shift reagent and high field (360 MHz) n.m.r. studies permitted the assignment of the proton resonances of alliacolide.

There are a number of biogenetic schemes that may lead to the formation of the carbon skeleton. $[1-^{13}\text{C}]$ - and $[1,2-^{13}\text{C}_2]$ -Acetate feeding experiments gave the enrichment and coupling pattern shown in figure 9, suggesting that the structure is made up of the combination of isoprene units indicated in figure 9. Significantly the carbonyl group of the lactone ring (C-13) is coupled to carbon-11, i.e. it must originate from the methyl group of mevalonic acid. In an isoprenoid chain, the methyl group of mevalonate remains a pendant group.




figure 8

Consequently the lactone ring may not have its origin in the cleavage of the isoprene chain. This result could even suggest that the isopropyl group rather than the gem-dimethyl group forms the starting point of the isoprene chain.




figure 9

The numbers of mevalonoid hydrogen atoms which were incorporated, were established by conventional ³H:¹⁴C ratio studies which are summarized in table 2.

Table 2 Incorporation of Mevalonates into Alliocolide (2)

Mevalonate	2- ³ H ₂ , 2- ¹⁴ C	4(R)-4- ³ H, 2- ¹⁴ C	5- ³ H ₂ , 2- ¹⁴ C
³ H: ¹⁴ C atom ratio in (2)	5.2:3	1.9:3	3.96:3
³ H: ¹⁴ C atom ratio in (5), obtained by degradation.	5.3:3	1.9:3	4:3
incorporation(%) into (2)	1.29	0.99	1.59

Two of the three centres (C-4 and C-9) which would be expected to be labelled by 4(R)-4-³H-mevalonate are fully substituted and hence to account for the incorporation of two labels, one pro-4(R)-mevalonoid hydrogen atom must be involved in a rearrangement. Secondly one of the centres (C-5) labelled by a 5-mevalonate is fully substituted and another (C-8) bears only one hydrogen atom. To account for the incorporation of the fourth (5-³H)-mevalonoid label, it must also be involved in a rearrangement. Simple dehydration to dehydroalliocolide (5) showed, surprisingly, that there was no mevalonoid label at C-11 whilst degradation to the hydroxy-ketone (6) established that C-12 bore two (2-³H)-mevalonoid hydrogen labels. However the remaining sites of label were determined using (2-³H₂)-acetate and the (2-³H₂)-, (4-³H₂)- and (5-³H₂)-mevalonates with deuterium n.m.r. spectroscopy. Although it was possible to assign the proton resonances of alliocolide, they were not sufficiently well resolved for satisfactory deuterium n.m.r. studies except in the case of the

(5- $^2\text{H}_2$)-MVA labelled material. However a satisfactory set of ^2H n.m.r. spectra were obtained from the degradation product, dehydroalliocolide (5). The results are summarized in figure 10. The hydrogen atom adjacent to the methyl group at C-1 is derived from the pro-4(R) position of mevalonate whilst a (5- ^2H)-mevalonoid label is attached to C-6. A small amount of scrambling of the (2- $^2\text{H}_2$)- and (4- $^2\text{H}_2$)-MVA labels was also apparent possibly due to degradation to acetate and re-synthesis. The absence of a mevalonoid label




figure 10

from C-11 is accounted for by the fact that dehydroalliocolide (5) is a very effective (32.7%) precursor of alliocolide (2). One possible biosynthetic scheme which would accommodate these results, but is not proven by them, is shown in figure 11. Such a scheme suggests an interesting role and origin of the epoxide oxygen atom.




figure 11

In conclusion, although detailed sequence studies remain to be done on both these fungal metabolites, I hope that the work illustrates the power of modern n.m.r. methods in biosynthetic problems.

Acknowledgements

I wish to acknowledge the experimental work of my student, Mr.A.P.W.Bradshaw, and the technical help of Mrs. N. Dransfield. Some preliminary experiments on dihydrobotrydial were carried out by Dr.R.Nyfelner. I am particularly indebted to Dr. I.H.Sadler of the University of Edinburgh for the determination of the high field ^1H , ^2H and ^{13}C n.m.r. spectra. I should also like to acknowledge the gift of a culture of Marasmius alliaceus and helpful discussions with Dr.T.G. Halsall and Dr.V.Thaller. The work was supported by the S.R.C.

REFERENCES

1. L.Ruzicka, Proc.Chem.Soc., 341 (1959).
2. for an application of computer methods to this see:- D.H.Smith and R.E.Carhart, Tetrahedron, 32, 2513 (1976); T.H.Varley, D.H.Smith and C.Djerassi, Tetrahedron, 34, 841 (1978).
3. D.Arigoni, Pure Appl.Chem., 41, 219 (1975); for a review see G.A.Cordell, Chem.Rev., 76, 425 (1976).
4. B.Achilladelis and J.R.Hanson, Chem. and Ind., 1643(1967); Phytochemistry, 7, 589 (1968); P.M.Adams and J.R.Hanson, JCS Perkin 1, 586 (1972); B.Achilladelis, P.M.Adams and J.R.Hanson, JCS Perkin 1, 1425 (1972); R.Evans, J.R.Hanson and T.Marten, JCS Perkin 1, 857 (1974); J.R.Hanson, T.Marten and M.Siverns, JCS Perkin 1, 1033 (1974); R.Evans and J.R.Hanson, JCS Perkin 1, 326 (1976); J.R.Hanson, T.Marten and R.Nyfelner, JCS Perkin 1, 876(1976); R.Evans, J.R.Hanson and T.Marten, JCS Perkin 1, 1212 (1976); R.Evans, J.R.Hanson and R.Nyfelner, JCS Perkin 1, 1214 (1976); J.R.Hanson and R.Nyfelner, JCS Perkin 1, 2471 (1976); JCS Chem. Comm., 72 (1976); A.P.W.Bradshaw, J.R.Hanson and M.Siverns, JCS Chem.Comm., 819(1977); JCS Chem.Comm., 303 (1978); A.P.W.Bradshaw and J.R.Hanson, JCS Chem.Comm., 924 (1979).
5. H-W. Fehlhaber, R.Geipal, H-J. Mercker, R.Tschesche and K.Welmar, Chem. Ber., 107, 1720 (1974); H.J.Linder and B.von Groasse, Chem.Ber., 107, 3332 (1974).
6. T.J.King, I.W.Farrell, T.G.Halsall and V.Thaller, JCS Chem.Comm., 727 (1977).
7. for reviews see:- T.J.Simpson, Chem.Soc.Rev., 4, 497(1975); M.J.Garson and J.Staunton, Chem.Soc. Rev., 8, 539 (1979).
8. B.Dockerill, J.R.Hanson and M.Siverns, Phytochemistry, 17, 427 (1978).
9. R.Croteau and W.D.Loomis, Phytochemistry, 11, 1055 (1972).
10. O.Cuevas and J.R.Hanson, Phytochemistry, 16, 1061 (1977); A.P.W.Bradshaw and J.R.Hanson, JCS Perkin 1, 741 (1980).